

COMMUNITY
URINALYSIS

SELF-REPORT PROJECT

METHAMPHETAMINE
WAS THE
ILLICIT DRUG
REPORTED TO BE USED
MOST OFTEN

46% OF PARTICIPANTS

AND WAS ALSO THE SUBSTANCE MOST
DETECTED IN THE URINALYSIS

IRI
M
H
N
MANITOBA
HARM REDUCTION
NETWORK



COMMUNITY
URINALYSIS

SELF-REPORT PROJECT

40%
OF RESPONDENTS
INDICATED THAT THEY USE
ILLEGAL/ILLICIT DRUGS
EVERY DAY

**11% OF RESPONDENTS USED NEEDLES OR
SYRINGES THAT HAD BEEN USED BY OTHERS**

**IRIM
IHI
IN**

MANITOBA
HARM REDUCTION
NETWORK



COMMUNITY
URINALYSIS

SELF-REPORT PROJECT

OF PARTICIPANTS THAT USE DEPRESSANTS:

17% PREFERRED TO USE HYDROMORPHONE

15% PREFERRED TO USE MORPHINE

OF PARTICIPANTS THAT USE STIMULANTS

37% PREFERRED TO USE METHAMPHETAMINE

26% PREFERRED TO USE COCAINE

SMOKING WAS THE PREFERRED METHOD
OF USE (61%) FOLLOWED BY INJECTING (21%)
AND SNORTING (11%)

